



# Measurement of $D^0$ decays to $D^0 \rightarrow K_L^0 \pi^0$ and $D^0 \rightarrow K_S^0 \pi^0$

Belle Analysis Meeting,  
11-12 Sept 2006

Manmohan Dash  
Virginia Tech



# In this presentation...

- Motivation for this analysis
- Selection procedure
- Recapitulation of previous studies
- Detailed Background study
- Priority issues and conclusion

This analysis intends to measure the asymmetry in  
 $D^0 \rightarrow K_L^0 \pi^0$  and  $D^0 \rightarrow K_S^0 \pi^0$ .

Physics Letters B 505(2001)94-106

- comes from interference in CF and DCS modes of  $D^0$  decay, expected up to 5%.
- constrains the strong phase in  $K^+ \pi^-$ ,  $K^- \pi^+$
- can be calibrated against  $D^0 \rightarrow (\bar{K}^0 \pi^-)_{K^*^-} \pi^+$
- measured at belle,  $0.06 \pm 0.05 \pm 0.05$

hep-ex/0107078, July 2001

- measured at cleo  $0.122 \pm 0.024 \pm 0.030$

hep-ex/0607068, July 2006

# Reconstruction and Event selection

- $\pi^0$  from mdst\_pi0,  $E_\gamma \geq 0.05$  GeV/c<sup>2</sup>
- $\pi^\pm$ , good charged
- $K_S$ , good Kshort
- $K_L$ , mdst\_klong,  $D^0 \rightarrow K_L \pi^0$  and  $D^0 \rightarrow (K_L \pi^-)_K^* \pi^+$  assumed
- $K^{*\pm}$  within 50 MeV of nominal mass
- $D^0$  within 100 MeV of nominal mass (for  $K_S$  modes only)
- $D^{*\pm}$  tags the signal,  $M_{D^{*\pm}} \leq 2.03$  GeV
- $\delta M = D^{*\pm}$  and  $D^0$  mass difference  
 $\delta M + 1.8645 \leq 2.03$  GeV  
(for  $K_S$  modes only)
- A  $K_L$  with ECL cluster energy in 0.15 to 0.3 GeV range (corresponding to minimum ionization energy) rejected
- $K^0$  flight angle wrt  $D^0$  boost ( $\theta_{DK}$ ),  $-0.95 \leq \cos(\theta_{DK}) \leq 0.2$  for all modes
- Invariant mass of  $(\pi^+ \pi^-) \leq 0.7$  GeV
- Reconstructed scaled momentum of  $D^*$ ,  $0.6 \leq x_p \leq 1.0$









# Comparison on same data

		This analysis	2001, Belle conf 0129
yield	$D^0 \rightarrow K_S \pi^0$	6739	$4715 \pm 91$
	$D^0 \rightarrow K_L \pi^0$	3943	$1839 \pm 101$
	$D^0 \rightarrow (K_S \pi^+) \pi^-$	3432	$2524 \pm 77$
	$D^0 \rightarrow (K_L \pi^+) \pi^-$	2617	$1119 \pm 84$
selection	$K_S$ selection	Good kshort	Tighter vertex
	$\pi^0 / \pi^\pm$ selection	$\pi^0$ from mdst_pi0, $E_\gamma \geq 0.05$ $\text{GeV}/c^2$ $\pi^\pm$ , good charged	Standard (?) procedure, 2001
fitting	Signal	Double Gaussian Mean of Gaussian fixed to $D^*$ nominal mass	Double Gaussian (KS modes), Single Gaussian (KL modes), means fixed
	Background	Threshold function	1st Poly x sqrt threshold 5

# On $23.6 \text{ fb}^{-1}$ from Experiment 7, 9, 11, 13

Upper: This Analysis,

Lower: Analysis from 2001



$D^0 \rightarrow K_S \pi^0$

$D^0 \rightarrow K_L \pi^0$

$D^0 \rightarrow K_S \pi^+ \pi^-$


$D^0 \rightarrow K_L \pi^+ \pi^-$

# The data-MC discrepancy !!, BAM Sept 2005

Data =  $32.37 \text{ fb}^{-1}$   
MC =  $\sim 2.7 \times \text{Data}$






Missing cont. data

Correct




# Detailed background study follows...

- MC truth study has been done by tagging specific decay modes at the generated level in the analysis code and vetoing or allowing them at the n-tuple level.
- Overall event composition study is shown in next page followed by detailed breakdown of background modes in the later slides.




$D^0 \rightarrow (K_L \pi^-) \pi^+$









Signal in Red  
Peaking background from  $\bar{K}^0/K^0 \rho^0$   
Background from other broad resonances  
Unknown peaking + rest of c cbar background  
uds background  
mixed + charged background

	yield	$\bar{K}R, R=\rho^0, \eta, \omega, f$ etc	$K^-\pi^+$	$K^*K^+$	$K^*K^-$
$D^0 \rightarrow K_S \pi^0$	13959. $\pm$ 133.06	91.49 $\pm$ 11.07 ( $\sim 0.6\%$ )	---	---	---
$D^0 \rightarrow K_L \pi^0$ KLM type	8957.6 $\pm$ 302.57	---	---	---	---
$D^0 \rightarrow K_L \pi^0$ ECL type	5313.6 $\pm$ 320.34	---	---	---	---




	yield	$\bar{K}R$ , $R = \rho^0$	$\bar{K}R$ , $R = \eta, \omega, f$ etc	$K^- \pi^+$	$K^* - K^+$	$K^* + K^-$	Still not known
PDG 2006 BF In %		$0.75 \pm 0.6 - 0.8$		$3.8 \pm 0.07$	$0.2 \pm 0.11$	$0.37 \pm 0.08$	
$D^0 \rightarrow (K_S \pi^-) \pi^+$	7937.6 $\pm/-$ 101.70	521.96 $\pm/-$ 26.49  <b>(5.76%)</b>	52.59 $\pm/-$ 8.23  <b>(0.58%)</b>	39.19 $\pm/-$ 7.48  <b>(0.43%)</b>	86.11 $\pm/-$ 9.56  <b>(0.83%)</b>	78.54 $\pm/-$ 8.91  <b>(0.87%)</b>	334.57 $\pm/-$ 56.56  <b>(3.70%)</b>
$D^0 \rightarrow (K_L \pi^-) \pi^+$ KLM type	4665.5 $\pm/-$ 121.76	354.22 $\pm/-$ 37.89  <b>(6.98%)</b>	55.88 $\pm/-$ 19.94  <b>(1.10%)</b>	---	---	---	---
$D^0 \rightarrow (K_S \pi^-) \pi^+$ ECL type	2334.3 $\pm/-$ 62.81	155.43 $\pm/-$ 16.05  <b>(3.58%)</b>	35.15 $\pm/-$ 8.08  <b>(0.81%)</b>	115.49 $\pm/-$ 26.22  <b>(2.66%)</b>	30.12 $\pm/-$ 46.03  <b>(0.69%)</b>	72.00 $\pm/-$ 21.56  <b>(1.66%)</b>	1595.8 $\pm/-$ 307.71  <b>(36.78%)</b>








# Peaking backgrounds in $D^0 \rightarrow (K_S \pi^-) \pi^+$



# Peaking backgrounds in $D^0 \rightarrow (K_L \pi^-) \pi^+$ , KLM type



# Peaking backgrounds in $D^0 \rightarrow (K_L \pi^-) \pi^+$ , ECL type



The unknown bump.  
If we discard all ECL  
KL then loss in  
signal efficiency:  
37.23% in signal  
mode.  
33.34% in  
calibration mode.


# Investigation for the unknown bump...

- Scanned ECL variables in calibration mode. Didn't find any strange behavior.
- Checked  $\pi^+ \pi^- \pi^0$  and  $\pi^+ \pi^- K_L$ . No peaking background from these sources.
- $K^- \pi^+, K^* - K^+, K^* + K^-$  don't account much for the bump.


# Conclusion.

- The peaking background yields will be estimated from MC. Uncertainty of PDG BF will introduce systematic error into asymmetry.
- The ECL in Calibration is suffering from most of the peaking backgrounds. It also has a strange huge bump with unknown source so far. We propose to discard the ECL solution for our estimation of asymmetry.


## uds background



## b bar background



# Multiple candidates



# Issues to be dealt on a priority basis.

- More MC statistics. Extend MC to exp 23. (~100/fb)  
Skim finished  $c\bar{c}$  MC. Extend the analysis to Exp 23 in MC.
- Belle Note started, will finish in 1 week.
- Bias and fit method systematics study from MC true # and fit yield.
- Asymmetry in data by smearing Pseudo KL direction to match KL direction resolution.
- No best candidate selection. Estimate error of this in asymmetry.
- Update to full data.
- Abstract submitted for DPF/JPS.